

# Clinical Appropriateness Guidelines: Arterial Ultrasound

## Appropriate Use Criteria

Effective Date: January 2, 2018

Proprietary

Date of Origin: 8/27/2015  
Last revised: 11/02/2017  
Last reviewed: 11/02/2017



8600 W Bryn Mawr Avenue  
South Tower - Suite 800  
Chicago, IL 60631  
P. 773.864.4600  
[www.aimspecialtyhealth.com](http://www.aimspecialtyhealth.com)

<b>Description and Application of the Guidelines</b> .....	<b>3</b>
<b>Arterial Ultrasound</b> .....	<b>4</b>
Duplex Ultrasound Imaging of the Extracranial Arteries.....	4
Duplex Ultrasound Imaging of the Aorta, Inferior Vena Cava and Iliac Vessels.....	7
Duplex Ultrasound Imaging of the Arteries of the Upper Extremities .....	11
Duplex Ultrasound Imaging of the Arteries of the Lower Extremities .....	14
Physiologic Testing for Peripheral Arterial Disease (PAD) of the Upper and Lower Extremities .....	17

# Description and Application of the Guidelines



AIM's Clinical Appropriateness Guidelines (hereinafter "AIM's Clinical Appropriateness Guidelines" or the "Guidelines") are designed to assist providers in making the most appropriate treatment decision for a specific clinical condition for an individual. As used by AIM, the Guidelines establish objective and evidence-based, where possible, criteria for medical necessity determinations. In the process, multiple functions are accomplished:

- To establish criteria for when services are medically necessary
- To assist the practitioner as an educational tool
- To encourage standardization of medical practice patterns
- To curtail the performance of inappropriate and/or duplicate services
- To advocate for patient safety concerns
- To enhance the quality of healthcare
- To promote the most efficient and cost-effective use of services

AIM's guideline development process complies with applicable accreditation standards, including the requirement that the Guidelines be developed with involvement from appropriate providers with current clinical expertise relevant to the Guidelines under review and be based on the most up to date clinical principles and best practices. Relevant citations are included in the "References" section attached to each Guideline. AIM reviews all of its Guidelines at least annually.

AIM makes its Guidelines publicly available on its website twenty-four hours a day, seven days a week. Copies of AIM's Clinical Appropriateness Guidelines are also available upon oral or written request. Although the Guidelines are publicly-available, AIM considers the Guidelines to be important, proprietary information of AIM, which cannot be sold, assigned, leased, licensed, reproduced or distributed without the written consent of AIM.

AIM applies objective and evidence-based criteria and takes individual circumstances and the local delivery system into account when determining the medical appropriateness of health care services. The AIM Guidelines are just guidelines for the provision of specialty health services. These criteria are designed to guide both providers and reviewers to the most appropriate services based on a patient's unique circumstances. In all cases, clinical judgment consistent with the standards of good medical practice should be used when applying the Guidelines. Guideline determinations are made based on the information provided at the time of the request. It is expected that medical necessity decisions may change as new information is provided or based on unique aspects of the patient's condition. The treating clinician has final authority and responsibility for treatment decisions regarding the care of the patient and for justifying and demonstrating the existence of medical necessity for the requested service. The Guidelines are not a substitute for the experience and judgment of a physician or other health care professionals. Any clinician seeking to apply or consult the Guidelines is expected to use independent medical judgment in the context of individual clinical circumstances to determine any patient's care or treatment.

The Guidelines do not address coverage, benefit or other plan specific issues. If requested by a health plan, AIM will review requests based on health plan medical policy/guidelines in lieu of AIM's Guidelines.

The Guidelines may also be used by the health plan or by AIM for purposes of provider education, or to review the medical necessity of services by any provider who has been notified of the need for medical necessity review, due to billing practices or claims that are not consistent with other providers in terms of frequency or some other manner.

---

CPT® (Current Procedural Terminology) is a registered trademark of the American Medical Association (AMA). CPT® five digit codes, nomenclature and other data are copyright by the American Medical Association. All Rights Reserved. AMA does not directly or indirectly practice medicine or dispense medical services. AMA assumes no liability for the data contained herein or not contained herein.

# Duplex Ultrasound Imaging of the Extracranial Arteries



## CPT Codes

- 93880..... Duplex scan of extracranial arteries; complete bilateral study
- 93882..... Duplex scan of extracranial arteries; unilateral or limited study

## Standard Anatomic Coverage

- Arteries of both the anterior (carotid) and posterior (vertebrobasilar) extracranial systems.

## Imaging Considerations

- This guideline does not supersede the enrollee's health plan medical policy specific to Duplex Imaging of the arteries of the upper and lower extremities
- Selection of the optimal diagnostic imaging for evaluation of extracranial arterial disease should be made within the context of other available modalities (which include CT angiography (CTA), Magnetic Resonance angiography (MRA) and contrast angiography) so that the resulting information facilitates patient management decisions and does not merely add a new layer of testing
- Duplicative testing or repeat imaging of the same anatomic area with same or similar technology may be subject to high-level review and may not be medically necessary unless there is a persistent diagnostic problem or there has been a change in clinical status (e.g. deterioration) or there is a medical intervention which warrants interval reassessment
- In some clinical situations, it may be appropriate to perform transcranial Doppler studies in addition to Duplex imaging. Transcranial Doppler studies are not subject to preauthorization and are therefore not addressed in this document
- For the purposes of this guideline symptoms are defined as follows:
  - Anterior symptoms (carotid vascular territory) include unilateral motor or sensory deficit, speech impairment, or amaurosis fugax
  - Posterior symptoms (vertebrobasilar territory) include vertigo, ataxia, diplopia, dysphagia, dysarthria
  - The terms cerebrovascular attack (CVA) and transient ischemic attack (TIA) do not apply specifically to either anterior or posterior circulation
- For the purposes of this guideline, severity of vascular stenosis is defined as follows:
  - Mild disease: <50% stenosis
  - Moderate disease: 50%-69% stenosis
  - Severe disease: 70%-99% stenosis
  - Total occlusion: 100% stenosis
- For the purposes of this guideline, the term "revascularization" should be taken to mean carotid endarterectomy, or stent implantation

# Duplex Imaging for Extracranial Arterial Disease

The following diagnostic indications for Duplex imaging for Extracranial Arterial Disease are accompanied by pre-test considerations as well as supporting clinical data and prerequisite information:

## Patients with suspected extracranial arterial disease

(any **one** of the following)

- New or worsening anterior or posterior neurological symptoms
  - This does not apply to patients with syncope or near-syncope
- Evaluation of syncope when cardiovascular causes (e.g. rhythm disturbance, valvular disease) have been excluded
- Hollenhorst plaque seen on retinal examination
- Evaluation for subclavian or vertebral steal syndrome in patients who develop lightheadedness or impaired vision in the setting of upper extremity exertion
- Evaluation for spontaneous carotid artery dissection in patients with a pulsatile neck mass
  - Iatrogenic or traumatic dissection is better evaluated with CTA or MRA

---

## Patients with established extracranial arterial disease who have not undergone revascularization

(any **one** of the following)

- New or worsening anterior or posterior neurological symptoms
  - This guideline does not apply to patients with syncope or near-syncope
- Evaluation of syncope when cardiovascular causes (e.g. Rhythm disturbance, valvular disease) have been excluded
- Surveillance studies every 6 months are appropriate for patients with severe (70-99%) carotid stenosis provided that the patient is a candidate for revascularization
- Annual surveillance studies (after the first year) are appropriate for patients with moderate (50-69%) stenosis provided that the patient is a candidate for revascularization

---

## Patients with established extracranial arterial disease who have undergone revascularization

(any **one** of the following)

- A baseline study (usually within 1 month following revascularization) is appropriate
- New or worsening neurological symptoms
- Two imaging studies (usually at about 6 and 12 months) are appropriate within the first year following revascularization
- Annual surveillance studies (after the first year) are appropriate
- Following an abnormal surveillance study revealing severe stenosis additional studies at six month intervals are appropriate provided that the patient is a candidate for repeat revascularization

## Reference / Literature Review

1. American College of Cardiology Foundation (ACCF); American College of Radiology (ACR); American Institute of Ultrasound in Medicine (AIUM); et al. ACCF/ACR/AIUM/ASE/ASN/ICAVL/SCAI/SCCT/SIR/SVM/SVS/SVU 2012 appropriate use criteria for peripheral vascular ultrasound and physiological testing part I: arterial ultrasound and physiological testing: a report of the American College of Cardiology Foundation appropriate use criteria task force, American College of Radiology, American Institute of Ultrasound in Medicine, American Society of Echocardiography, American Society of Nephrology, Intersocietal Commission for the Accreditation of Vascular Laboratories, Society for Cardiovascular Angiography and Interventions, Society of Cardiovascular Computed Tomography, Society for Interventional Radiology, Society for Vascular Medicine, Society for Vascular Surgery, [corrected] and Society for Vascular Ultrasound. *J Am Coll Cardiol.* 2012;60(3):242-276.
2. Brott TG, Halperin JL, Abbara S, et al. 2011 ASA/ACCF/AHA/AANN/AANS/ACR/ASNR/CNS/SAIP/SCAI/SIR/SNIS/SVM/SVS guideline on the management of patients with extracranial carotid and vertebral artery disease: a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines, and the American Stroke Association, American Association of Neuroscience Nurses, American Association of Neurological Surgeons, American College of Radiology, American Society of Neuroradiology, Congress of Neurological Surgeons, Society of Atherosclerosis Imaging and Prevention, Society for Cardiovascular Angiography and Interventions, Society of Interventional Radiology, Society of NeuroInterventional Surgery, Society for Vascular Medicine, and Society for Vascular Surgery. *J Am Coll Cardiol.* 2011;57(8):e16-e94.
3. Norgren L, Hiatt WR, Dormandy JA, et al. Inter-Society Consensus for the Management of Peripheral Arterial Disease (TASC II). *Eur J Vasc Endovasc Surg.* 2007;33 Suppl 1:S1-S75.
4. LeFevre ML, U.S. Preventive Services Task Force. Screening for asymptomatic carotid artery stenosis: U.S. Preventive Services Task Force recommendation. *Ann Intern Med.* doi:10.7326/M14-1333
5. Quershi AJ, Alexandrov AV, Tegeler CH, et al. Guidelines for Screening of Extracranial Carotid Artery Disease: a Statement for Healthcare Professionals from the Multidisciplinary Practice Guidelines Committee of the American Society of Neuroimaging. *J Neuroimaging.* 2007;17:19-47

# Duplex Ultrasound Imaging of the Aorta, Inferior Vena Cava and Iliac Vessels



## CPT Codes

- 93978..... Duplex scan of aorta, inferior vena cava, iliac vasculature, or bypass grafts; complete study
- 93979..... Duplex scan of aorta, inferior vena cava, iliac vasculature, or bypass grafts; unilateral or limited study

## Standard Anatomic Coverage

- Abdominal aorta, inferior vena cava (IVC), iliac vasculature and bypass grafts involving these vessels.

## Imaging Considerations

- This guideline does not supersede the enrollee's health plan medical policy specific to Duplex Imaging of the aorta, IVC, iliac vasculature and associated bypass grafts
- Selection of the optimal diagnostic imaging for evaluation of disease of the abdominal aorta, IVC or iliac vasculature should be made within the context of other available modalities (which include CT angiography (CTA), Magnetic Resonance angiography (MRA) and contrast angiography) so that the resulting information facilitates patient management decisions and does not merely add a new layer of testing
- Duplicative testing or repeat imaging of the same anatomic area with same or similar technology may be subject to high-level review and may not be medically necessary unless there is a persistent diagnostic problem or there has been a change in clinical status (e.g. deterioration) or there is a medical intervention which warrants interval reassessment
- When screening for abdominal aortic aneurysm (AAA), a negative study should not be followed by additional interval screening studies
- Screening for AAA is only appropriate in patients who are candidates for (and are willing to undergo) repair procedures
- For the purposes of this guideline the term "repair" when used in discussion of aortoiliac disease should be taken to mean any of the following; open surgical repair of AAA, aorto-iliac endograft placement for management of aorto-iliac aneurysm, aortic or iliac stent placement, or surgical bypass procedures
- The periodic surveillance guidelines for patients with AAA who have not undergone repair are based on maximum external aortic diameter
- For the purposes of this guideline, symptoms are defined as follows:
  - Claudication is defined as muscle fatigue, cramping, or pain that reproducibly begins during exercise and that promptly resolves with rest
  - Rest pain is similar to the pain of claudication but it occurs at rest in a patient with an established diagnosis of PAD or with physical examination evidence of advanced PAD such as markedly diminished pulses, gangrene or ulceration
  - Atypical symptoms describe limb pains other than rest pain or claudication

## Duplex Imaging of the Aorta and Iliac Arteries

The following diagnostic indications for Duplex imaging for of the aorta, IVC and iliac vessels are accompanied by pre-test considerations as well as supporting clinical data and prerequisite information:

### **Asymptomatic patients with neither signs nor symptoms of disease (screening studies)** (any one of the following)

- A single screening study for AAA is appropriate in male patients aged 60-75 years who have a first degree relative with AAA
- A single screening study for AAA is appropriate in male patients aged 60-75 years who are current or former smokers
- A single screening study for AAA is appropriate for female patients aged 60-75 who have a first degree relative with AAA and are current or former smokers

---

## Patients with suspected aorto-iliac disease who have abnormal signs or symptoms

(any **one** of the following)

- Patient with a pulsatile abdominal mass
- Patient with suspected or established femoral or popliteal artery aneurysm
- Patient with a thoracic aortic aneurysm
- Abnormal abdominal X-ray suggestive of AAA
- Patient with decreased or absent femoral pulse
- Patients with lower extremity claudication
- Patients with abnormal physiological testing suggesting aorto-iliac disease
- Patient with abdominal or femoral bruit
- Patients with evidence of atheroembolic disease of the lower extremities (ischemic or discolored toes, livedo reticularis etc.)

---

## Patients with established aorto-iliac disease who have not undergone repair

Patients with an established diagnosis of AAA may undergo imaging in any **one** of the following situations:

- New or worsening symptoms or signs of aorto-iliac disease
- AAA greater than or equal to 4.5 cm may undergo surveillance imaging as frequently as every six (6) months
  - In patients with aneurysms greater than or equal to 5.5 cm, considerations should be given to repair unless there are factors which significantly increase procedural risk
- AAA greater than or equal to 3.5 cm and less than 4.5 cm may undergo surveillance imaging at six (6) monthly intervals in the first year following diagnosis and annually thereafter
- AAA greater than or equal to 3.0 cm and less than 3.5 cm may undergo surveillance imaging one year after diagnosis and then every three (3) years thereafter
- Iliac aneurysms greater than or equal to 3.0 cm may undergo surveillance imaging as frequently as every six (6) months
  - In patients with aneurysms greater than or equal to 3.5 cm, considerations should be given to repair unless there are factors which significantly increase procedural risk
- Iliac aneurysm less than 3.0 cm may undergo surveillance imaging annually

---

## Patients with established aorto-iliac disease who have undergone repair

Patients who have undergone aorto-iliac repair (as defined in imaging considerations above) may undergo imaging in any **one** of the following situations:

- New or worsening symptoms or signs of aorto-iliac disease
- Baseline study after aortic or iliac stent placement
- Baseline study after aortic or iliac endograft
  - Some providers will use CT angiography rather than ultrasound in this situation
- Follow-up surveillance study at 6 monthly intervals after aortic or iliac endograft when endograft leak or increasing residual aneurysm sac size was noted on the preceding study
  - Some providers will use CT angiography rather than ultrasound in this situation
- Follow-up surveillance study at yearly intervals after aortic or iliac endograft when there was no evidence of endograft leak or increasing residual aneurysm sac size on the preceding study
- Baseline study of surgical bypass grafting involving the aorto-iliac vessels
- Follow-up surveillance study at 6-12 months following surgical bypass grafting involving the aorto-iliac vessels
- Annual follow-up surveillance study starting 1 year after surgical bypass grafting involving the aorto-iliac vessels

## Duplex Imaging of the IVC and Iliac Veins

The following diagnostic indications for Duplex imaging for of the IVC and iliac veins are accompanied by pre-test considerations as well as supporting clinical data and prerequisite information:

### **Duplex imaging of the IVC and iliac vessels is appropriate in any one of the following situations:**

- Suspected or established IVC or iliac vein thrombus
- Suspected or established IVC or iliac vein mass
- Suspected or established external compression of the IVC or iliac veins
- To establish patency of the IVC in a patient with an IVC filter
- To evaluate tumor extension into the IVC in patients with renal or adrenal tumors
- To assist in evaluation of volume status in patients with unexplained hypotension (not usually performed in the outpatient setting)

## Reference/Literature Review

1. American College of Cardiology Foundation (ACCF); American College of Radiology (ACR); American Institute of Ultrasound in Medicine (AIUM); et al. ACCF/ACR/AIUM/ASE/ASN/ICAVL/SCAI/SCCT/SIR/SVM/SVS/SVU 2012 appropriate use criteria for peripheral vascular ultrasound and physiological testing part I: arterial ultrasound and physiological testing: a report of the American College of Cardiology Foundation appropriate use criteria task force, American College of Radiology, American Institute of Ultrasound in Medicine, American Society of Echocardiography, American Society of Nephrology, Intersocietal Commission for the Accreditation of Vascular Laboratories, Society for Cardiovascular Angiography and Interventions, Society of Cardiovascular Computed Tomography, Society for Interventional Radiology, Society for Vascular Medicine, Society for Vascular Surgery, [corrected] and Society for Vascular Ultrasound. *J Am Coll Cardiol*. 2012;60(3):242-276.
2. American College of Cardiology Foundation, American Heart Association, Inc. ACCF/AHA Pocket Guideline: Management of Patients With Peripheral Artery Disease (Lower Extremity, Renal, Mesenteric, and Abdominal Aortic). ACCF/AHA; November 2011.
3. Centers for Medicare and Medicaid Services. Chapter 18 – Preventive and Screening Services. Medicare Claims Processing Manual. Pub. 100-04. Section 110 – Ultrasound Screening for Abdominal Aortic Aneurysm (AAA). (Rev. 3763, Issued: 04-28-17) <http://www.cms.gov/Regulations-and-Guidance/Guidance/Manuals/Downloads/clm104c18.pdf>.
4. Chaikof EL, Brewster DC, Dalman RL, et al. The care of patients with an abdominal aortic aneurysm: The Society for Vascular Surgery Practice Guidelines. *J Vasc Surg*. 2009;50 Suppl:S2-S49.
5. Ferket BS, Grootenboer N, Colkesen EB, et al. Systematic review of guidelines on abdominal aortic aneurysm Screening. *J Vasc Surg*. 2012;55(5):1296-1304.
6. Hirsch AT, Haskal ZJ, Hertzner NR, et al. ACC/AHA 2005 Practice Guidelines for the management of patients with peripheral arterial disease (lower extremity, renal, mesenteric, and abdominal aortic): a collaborative report from the American Association for Vascular Surgery/Society for Vascular Surgery, Society for Cardiovascular Angiography and Interventions, Society for Vascular Medicine and Biology, Society of Interventional Radiology, and the ACC/AHA Task Force on Practice Guidelines (Writing Committee to Develop Guidelines for the Management of Patients With Peripheral Arterial Disease): endorsed by the American Association of Cardiovascular and Pulmonary Rehabilitation; National Heart, Lung, and Blood Institute; Society for Vascular Nursing; TransAtlantic Inter-Society Consensus; and Vascular Disease Foundation. *Circulation*. 2006;113(11):e463-e654.
7. Kasirajan V, Hertzner NR, Beven EG, O'Hara PJ, Krajewski LP, Sullivan TM. Management of isolated common iliac artery aneurysms. *Cardiovasc Surg*. 1998;6(2):171.
8. Krupski WC, Selzman CH, Florida R, Strecker PK, Nehler MR, Whitehill TA. Contemporary management of isolated iliac aneurysms. *J Vasc Surg*. 1998;28(1):1.
9. LeFevre ML; U.S. Preventive Services Task Force. Screening for Abdominal Aortic Aneurysm: U.S. Preventive Services Task Force Recommendation Statement. *Ann Intern Med*. 2014 Aug 19;161(4):281-290.
10. Mastracci TM, Cina CS; Canadian Society for Vascular Surgery. Screening for abdominal aortic aneurysm in Canada: review and position statement of the Canadian Society for Vascular Surgery. *J Vasc Surg* 2007;45:1268-1276.
11. Norgren L, Hiatt WR, Dormandy JA, et al. Inter-society consensus for the management of peripheral arterial disease (TASC II). *Eur J Vasc Endovasc Surg*. 2007;33 Suppl 1:S1-S75.
12. RESCAN Collaborators, Bown MJ, Sweeting MJ, Brown LC, Powell JT, Thompson SG. Surveillance intervals for small abdominal aortic aneurysms: a meta-analysis. *JAMA*. 2013 Feb 27;309(8):806-813.
13. Richardson JW, Greenfield LJ. Natural history and management of iliac aneurysms. *J Vasc Surg*. 1988;8(2):165.
14. Rooke TW, Hirsch AT, Misra S, et al. 2011 ACCF/AHA Focused Update of the Guideline for the Management of Patients With Peripheral Artery Disease (updating the 2005 guideline): a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines. *J Am Coll Cardiol*. 2011;58(19):2020-2045.
15. Rooke TW, Hirsch AT, Misra S, et al.; American College of Cardiology Foundation Task Force; American Heart Association Task Force. Management of patients with peripheral artery disease (compilation of 2005 and 2011 ACCF/AHA Guideline Recommendations): a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines. *J Am Coll Cardiol*. 2013;61(14):1555-1570.
16. Santilli SM, Wernsing SE, Lee ES. Expansion rates and outcomes for iliac artery aneurysms. *J Vasc Surg*. 2000;31(1 Pt 1):114.
17. Stavropoulos SW, Charagundla SR. Imaging techniques for detection and management of endoleaks after endovascular aortic aneurysm repair. *Radiology*. 2007 Jun;243(3):641-655.
18. Upchurch GR Jr, Schaub TA. Abdominal aortic aneurysm. *Am Fam Physician*. 2006 Apr 1;73(7):1198-1204.

# Duplex Ultrasound Imaging of the Arteries of the Upper Extremities



## CPT Codes

- 93930..... Duplex scan of upper extremity arteries or arterial bypass grafts; complete bilateral study
- 93931..... Duplex scan of upper extremity arteries or arterial bypass grafts; unilateral or limited study

## Standard Anatomic Coverage

- Arteries of the upper extremities

## Imaging Considerations

- This guideline does not supersede the enrollee's health plan medical policy specific to Duplex Imaging of the arteries of the upper extremities
- This guideline does not address physiological imaging of the upper extremities (CPT codes 93922-93924) which are covered in a separate guideline document
- Selection of the optimal diagnostic imaging for evaluation of peripheral arterial disease should be made within the context of other available modalities (which include Duplex Vascular imaging studies, CT angiography (CTA), Magnetic Resonance angiography (MRA) and contrast angiography) so that the resulting information facilitates patient management decisions and does not merely add a new layer of testing
- Duplicative testing or repeat imaging of the same anatomic area with same or similar technology may be subject to high-level review and may not be medically necessary unless there is a persistent diagnostic problem or there has been a change in clinical status (e.g. deterioration) or there is a medical intervention which warrants interval reassessment
- It is assumed that patients who have cardiovascular disease (established coronary, cerebrovascular disease) and patients at high risk of cardiovascular disease (including patients with diabetes mellitus and chronic kidney disease) will be treated with optimal medical therapy. Therefore, screening for asymptomatic peripheral artery disease (PAD) in these populations is unlikely to change management
- Evidence is lacking that treatment of asymptomatic PAD delays the onset of symptomatic PAD
- In general (exceptions noted below), physiological imaging should be the initial imaging approach to the evaluation of PAD. Duplex imaging should be reserved for situations in which physiological studies are inconclusive or when physiological studies are abnormal, the patient had failed conservative therapy and the patient is being evaluated for revascularization
- It is conventional to report ABI measurements as follows: Noncompressible values defined as greater than 1.40, normal values 1.00 to 1.40, borderline 0.91 to 0.99, and abnormal 0.90 or less
- For the purposes of this guideline symptoms are defined as follows:
  - Claudication is defined as muscle fatigue, cramping, or pain that reproducibly begins during exercise and that promptly resolves with rest
  - Rest pain is similar to the pain of claudication but it occurs at rest in a patient with an established diagnosis of PAD or with physical examination evidence of advanced PAD such as markedly diminished pulses, gangrene or ulceration
  - Atypical symptoms describe limb pains other than rest pain or claudication

# Duplex Imaging for Peripheral Arterial Disease of the Upper Extremities

The following diagnostic indications for Duplex imaging for Peripheral Arterial Disease of the Upper Extremities are accompanied by pre-test considerations as well as supporting clinical data and prerequisite information:

## Asymptomatic patients with suspected PAD

- Screening for asymptomatic PAD has not been shown to affect outcomes and is therefore not considered medically necessary

---

## Symptomatic patients with suspected PAD (see definition of symptoms under Imaging Considerations above)

(any **one** of the following)

- Duplex imaging is appropriate for patients with atypical symptoms who have inconclusive physiological testing
- Duplex imaging is appropriate for patients with claudication who have normal or inconclusive physiological testing
- Patients with resting ischemic pain
- Patients with evidence of atheroembolic disease of the upper extremities (ischemic or discolored fingers, livedo reticularis etc.)

---

## Patients with established PAD

(any **one** of the following)

- Duplex imaging is appropriate for patients with atypical symptoms who have inconclusive physiological testing
- Duplex imaging is appropriate for patients with claudication who have normal or inconclusive physiological testing
- Patients with resting ischemic pain
- Patients with evidence of atheroembolic disease of the upper extremities (ischemic or discolored fingers, livedo reticularis etc.)
- Patients who have persistent claudication despite a trial of conservative therapy who are being evaluated for revascularization
- A routine baseline study is appropriate for patients who have undergone revascularization (percutaneous or surgical)
- Duplex imaging is appropriate for patients who have undergone revascularization when surveillance (no new or worsening symptoms) physiological testing is inconclusive
- A follow-up surveillance (no new or worsening symptoms) study at 6-12 months following surgical revascularization is appropriate. Note that this guideline is not applicable for surveillance following percutaneous revascularization (angioplasty, stent placement etc.)
- An annual follow-up surveillance (no new or worsening symptoms) study starting 1 year after revascularization is appropriate for patients who have undergone surgical revascularization. Note that this guideline is not applicable for surveillance following percutaneous revascularization (angioplasty, stent placement etc.)

---

## Patients who have had procedures requiring arterial access

Duplex imaging is appropriate for evaluation of vascular access complications when a patient who has had vascular access has any **one** of the following

- A pulsatile mass
- A bruit or thrill at the vascular access site
- A significant (more than would be expected for the procedure performed) hematoma at the vascular access site
- Severe pain (more than would be expected for the procedure performed) at the vascular access site
- Patients with evidence of atheroembolic disease of the upper extremities (ischemic or discolored fingers, livedo reticularis etc.)

---

## Miscellaneous indications for duplex imaging

(any **one** of the following)

- Following limb trauma when there is suspicion of vascular injury
- To assess the suitability of upper extremity arteries for use as bypass conduits prior to CABG
- For evaluation of suspected positional arterial obstruction (e.g. thoracic outlet syndrome)

## Reference/Literature Review

1. American College of Cardiology Foundation (ACCF); American College of Radiology (ACR); American Institute of Ultrasound in Medicine (AIUM); et al. ACCF/ACR/AIUM/ASE/ASN/ICAVL/SCAI/SCCT/SIR/SVM/SVS/SVU 2012 appropriate use criteria for peripheral vascular ultrasound and physiological testing part I: arterial ultrasound and physiological testing: a report of the American College of Cardiology Foundation appropriate use criteria task force, American College of Radiology, American Institute of Ultrasound in Medicine, American Society of Echocardiography, American Society of Nephrology, Intersocietal Commission for the Accreditation of Vascular Laboratories, Society for Cardiovascular Angiography and Interventions, Society of Cardiovascular Computed Tomography, Society for Interventional Radiology, Society for Vascular Medicine, Society for Vascular Surgery, [corrected] and Society for Vascular Ultrasound. *J Am Coll Cardiol*. 2012;60(3):242-276.
2. American College of Cardiology Foundation, American Heart Association, Inc. *ACCF/AHA Pocket Guideline: Management of Patients With Peripheral Artery Disease (Lower Extremity, Renal, Mesenteric, and Abdominal Aortic)*. ACCF/AHA; November 2011.
3. Brott TG, Halperin JL, Abbara S, et al. 2011 ASA/ACCF/AHA/AANN/AANS/ACR/ASNR/CNS/SAIP/SCAI/SIR/SNIS/SVM/SVS guideline on the management of patients with extracranial carotid and vertebral artery disease: a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines, and the American Stroke Association, American Association of Neuroscience Nurses, American Association of Neurological Surgeons, American College of Radiology, American Society of Neuroradiology, Congress of Neurological Surgeons, Society of Atherosclerosis Imaging and Prevention, Society for Cardiovascular Angiography and Interventions, Society of Interventional Radiology, Society of NeuroInterventional Surgery, Society for Vascular Medicine, and Society for Vascular Surgery. *J Am Coll Cardiol*. 2011;57(8):e16-e94.
4. Hirsch AT, Haskal ZJ, Hertzner NR, et al. ACC/AHA 2005 Practice Guidelines for the management of patients with peripheral arterial disease (lower extremity, renal, mesenteric, and abdominal aortic): a collaborative report from the American Association for Vascular Surgery/Society for Vascular Surgery, Society for Cardiovascular Angiography and Interventions, Society for Vascular Medicine and Biology, Society of Interventional Radiology, and the ACC/AHA Task Force on Practice Guidelines (Writing Committee to Develop Guidelines for the Management of Patients With Peripheral Arterial Disease): endorsed by the American Association of Cardiovascular and Pulmonary Rehabilitation; National Heart, Lung, and Blood Institute; Society for Vascular Nursing; TransAtlantic Inter-Society Consensus; and Vascular Disease Foundation. *Circulation*. 2006;113(11):e463-e654.
5. Moyer VA; U.S. Preventive Services Task Force. Screening for peripheral artery disease and cardiovascular disease risk assessment with the ankle-brachial index in adults: U.S. Preventive Services Task Force recommendation statement. *Ann Intern Med*. 2013;159(5):342-348.
6. Norgren L, Hiatt WR, Dormandy JA, et al. Inter-Society Consensus for the Management of Peripheral Arterial Disease (TASC II). *Eur J Vasc Endovasc Surg*. 2007;33 Suppl 1:S1-S75.
7. Rooke TW, Hirsch AT, Misra S, et al. 2011 ACCF/AHA Focused Update of the Guideline for the Management of Patients With Peripheral Artery Disease (updating the 2005 guideline): a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines. *J Am Coll Cardiol*. 2011;58(19):2020-2045.
8. Rooke TW, Hirsch AT, Misra S, et al.; American College of Cardiology Foundation Task Force; American Heart Association Task Force. Management of patients with peripheral artery disease (compilation of 2005 and 2011 ACCF/AHA Guideline Recommendations): a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines. *J Am Coll Cardiol*. 2013;61(14):1555-1570.

# Duplex Ultrasound Imaging of the Arteries of the Lower Extremities



## CPT Codes

93925..... Duplex scan of lower extremity arteries or arterial bypass grafts; complete bilateral study

93926..... Duplex scan of lower extremity arteries or arterial bypass grafts; unilateral or limited study

## Standard Anatomic Coverage

- Arteries of the lower extremities

## Imaging Considerations

- This guideline does not supersede the enrollee's health plan medical policy specific to Duplex Imaging of the arteries of the lower extremities
- This guideline does not address physiological imaging of the lower extremities (CPT codes 93922-93924) which are covered in a separate guideline document
- Selection of the optimal diagnostic imaging for evaluation of peripheral arterial disease should be made within the context of other available modalities (which include Duplex Vascular imaging studies, CT angiography (CTA), Magnetic Resonance angiography (MRA) and contrast angiography) so that the resulting information facilitates patient management decisions and does not merely add a new layer of testing
- Duplicative testing or repeat imaging of the same anatomic area with same or similar technology may be subject to high-level review and may not be medically necessary unless there is a persistent diagnostic problem or there has been a change in clinical status (e.g. deterioration) or there is a medical intervention which warrants interval reassessment
- It is assumed that patients who have cardiovascular disease (established coronary, cerebrovascular disease) and patients at high risk of cardiovascular disease (including patients with diabetes mellitus and chronic kidney disease) will be treated with optimal medical therapy. Therefore, screening for asymptomatic peripheral artery disease (PAD) in these populations is unlikely to change management
- Evidence is lacking that treatment of asymptomatic PAD delays the onset of symptomatic PAD
- In general (exceptions noted below), physiological imaging should be the initial imaging approach to the evaluation of PAD. Duplex imaging should be reserved for situations in which physiological studies are inconclusive or when physiological studies are abnormal, the patient had failed conservative therapy and the patient is being evaluated for revascularization
- It is conventional to report ABI measurements as follows: Noncompressible/inconclusive values defined as greater than 1.40, normal values 1.00 to 1.40, borderline 0.91 to 0.99, and abnormal 0.90 or less
- For the purposes of this guideline symptoms are defined as follows:
  - Claudication is defined as muscle fatigue, cramping, or pain that reproducibly begins during exercise and that promptly resolves with rest
  - Rest pain is similar to the pain of claudication but it occurs at rest in a patient with an established diagnosis of PAD or with physical examination evidence of advanced PAD such as markedly diminished pulses, gangrene or ulceration
  - Atypical symptoms describe limb pains other than rest pain or claudication

## Duplex Imaging for Peripheral Arterial Disease of the Lower Extremities

The following diagnostic indications for Duplex imaging for Peripheral Arterial Disease of the Lower Extremities are accompanied by pre-test considerations as well as supporting clinical data and prerequisite information:

### Asymptomatic patients with suspected PAD

- Screening for asymptomatic PAD has not been shown to affect outcomes and is therefore not considered medically necessary

---

## Symptomatic patients with suspected PAD (see definition of symptoms under Imaging

### Considerations above)

#### (any one of the following)

- Duplex imaging is appropriate for patients with atypical symptoms who have inconclusive physiological testing (e.g. ABI >1.40)
- Duplex imaging is appropriate for patients with claudication who have normal, borderline, or inconclusive physiological testing (ABI > 0.90)
- Patients with resting ischemic pain
- Patients with evidence of atheroembolic disease of the lower extremities (ischemic or discolored toes, livedo reticularis etc.)

---

## Patients with established PAD

#### (any one of the following)

- Duplex imaging is appropriate for patients with atypical symptoms who have inconclusive physiological testing (e.g. ABI >1.40)
- Duplex imaging is appropriate for patients with claudication who have normal, borderline, or inconclusive physiological testing (ABI > 0.90)
- Patients with resting ischemic pain
- Patients with evidence of atheroembolic disease of the lower extremities (ischemic or discolored toes, livedo reticularis etc.)
- Patients who have persistent claudication despite a trial of conservative therapy who are being evaluated for revascularization
- A routine baseline study is appropriate for patients who have undergone revascularization (percutaneous or surgical)
- Duplex imaging is appropriate for patients who have undergone revascularization when surveillance (no new or worsening symptoms) physiological testing is inconclusive (ABI >1.40), borderline (ABI 0.91 – 0.99) or abnormal (ABI < or = 0.90)
- A follow-up surveillance (no new or worsening symptoms) study at 6-12 months following surgical revascularization is appropriate. Note that this guideline is not applicable for surveillance following percutaneous revascularization (angioplasty, stent placement etc.)
- An annual follow-up surveillance (no new or worsening symptoms) study starting 1 year after revascularization is appropriate for patients who have undergone surgical revascularization. Note that this guideline is not applicable for surveillance following percutaneous revascularization (angioplasty, stent placement etc.)

---

## Patients who have had procedures requiring arterial access

Duplex imaging is appropriate for evaluation of vascular access complications when a patient who has had vascular access has **any one of the following**

- A pulsatile mass
- A bruit or thrill at the vascular access site
- A significant (more than would be expected for the procedure performed) hematoma at the vascular access site
- Severe pain (more than would be expected for the procedure performed) at the vascular access site
- Patients with evidence of atheroembolic disease of the lower extremities (ischemic or discolored toes, livedo reticularis etc.)

---

## Miscellaneous indications for duplex imaging

#### (any one of the following)

- Following limb trauma when there is suspicion of vascular injury
- For evaluation of suspected positional arterial obstruction

## Reference/Literature Review

1. American College of Cardiology Foundation (ACCF); American College of Radiology (ACR); American Institute of Ultrasound in Medicine (AIUM); et al. ACCF/ACR/AIUM/ASE/ASN/ICAVL/SCAI/SCCT/SIR/SVM/SVS/SVU 2012 appropriate use criteria for peripheral vascular ultrasound and physiological testing part I: arterial ultrasound and physiological testing: a report of the American College of Cardiology Foundation appropriate use criteria task force, American College of Radiology, American Institute of Ultrasound in Medicine, American Society of Echocardiography, American Society of Nephrology, Intersocietal Commission for the Accreditation of Vascular Laboratories, Society for Cardiovascular Angiography and Interventions, Society of Cardiovascular Computed Tomography, Society for Interventional Radiology, Society for Vascular Medicine, Society for Vascular Surgery, [corrected] and Society for Vascular Ultrasound. *J Am Coll Cardiol*. 2012;60(3):242-276.
2. American College of Cardiology Foundation, American Heart Association, Inc. *ACCF/AHA Pocket Guideline: Management of Patients With Peripheral Artery Disease (Lower Extremity, Renal, Mesenteric, and Abdominal Aortic)*. ACCF/AHA; November 2011.
3. Brott TG, Halperin JL, Abbara S, et al. 2011 ASA/ACCF/AHA/AANN/AANS/ACR/ASNR/CNS/SAIP/SCAI/SIR/SNIS/SVM/SVS guideline on the management of patients with extracranial carotid and vertebral artery disease: a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines, and the American Stroke Association, American Association of Neuroscience Nurses, American Association of Neurological Surgeons, American College of Radiology, American Society of Neuroradiology, Congress of Neurological Surgeons, Society of Atherosclerosis Imaging and Prevention, Society for Cardiovascular Angiography and Interventions, Society of Interventional Radiology, Society of NeuroInterventional Surgery, Society for Vascular Medicine, and Society for Vascular Surgery. *J Am Coll Cardiol*. 2011;57(8):e16-e94.
4. Hirsch AT, Haskal ZJ, Hertzner NR, et al. ACC/AHA 2005 Practice Guidelines for the management of patients with peripheral arterial disease (lower extremity, renal, mesenteric, and abdominal aortic): a collaborative report from the American Association for Vascular Surgery/Society for Vascular Surgery, Society for Cardiovascular Angiography and Interventions, Society for Vascular Medicine and Biology, Society of Interventional Radiology, and the ACC/AHA Task Force on Practice Guidelines (Writing Committee to Develop Guidelines for the Management of Patients With Peripheral Arterial Disease): endorsed by the American Association of Cardiovascular and Pulmonary Rehabilitation; National Heart, Lung, and Blood Institute; Society for Vascular Nursing; TransAtlantic Inter-Society Consensus; and Vascular Disease Foundation. *Circulation*. 2006;113(11):e463-e654.
5. Moyer VA; U.S. Preventive Services Task Force. Screening for peripheral artery disease and cardiovascular disease risk assessment with the ankle-brachial index in adults: U.S. Preventive Services Task Force recommendation statement. *Ann Intern Med*. 2013;159(5):342-348.
6. Norgren L, Hiatt WR, Dormandy JA, et al. Inter-Society Consensus for the Management of Peripheral Arterial Disease (TASC II). *Eur J Vasc Endovasc Surg*. 2007;33 Suppl 1:S1-S75.
7. Rooke TW, Hirsch AT, Misra S, et al. 2011 ACCF/AHA Focused Update of the Guideline for the Management of Patients With Peripheral Artery Disease (updating the 2005 guideline): a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines. *J Am Coll Cardiol*. 2011;58(19):2020-2045.
8. Rooke TW, Hirsch AT, Misra S, et al.; American College of Cardiology Foundation Task Force; American Heart Association Task Force. Management of patients with peripheral artery disease (compilation of 2005 and 2011 ACCF/AHA Guideline Recommendations): a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines. *J Am Coll Cardiol*. 2013;61(14):1555-1570.

## CPT Codes

- 93922..... Limited bilateral noninvasive physiologic studies of upper or lower extremity arteries, (e.g., for lower extremity: ankle/brachial indices at distal posterior tibial and anterior tibial/dorsalis pedis arteries plus bidirectional, Doppler waveform recording and analysis at 1-2 levels, or ankle/brachial indices at distal posterior tibial and anterior tibial/dorsalis pedis arteries plus volume plethysmography at 1-2 levels, or ankle/brachial indices at distal posterior tibial and anterior tibial/dorsalis pedis arteries with, transcutaneous oxygen tension measurement at 1-2 levels)
- 93923..... Complete bilateral noninvasive physiologic studies of upper or lower extremity arteries, 3 or more levels (e.g., for lower extremity: ankle/brachial indices at distal posterior tibial and anterior tibial/dorsalis pedis arteries plus segmental blood pressure measurements with bidirectional Doppler waveform recording and analysis, at 3 or more levels, or ankle/brachial indices at distal posterior tibial and anterior tibial/dorsalis pedis arteries plus segmental volume plethysmography at 3 or more levels, or ankle/brachial indices at distal posterior tibial and anterior tibial/dorsalis pedis arteries plus segmental transcutaneous oxygen tension measurements at 3 or more levels), or single level study with provocative functional maneuvers (e.g., measurements with postural provocative tests, or measurements with reactive hyperemia)
- 93924..... Noninvasive physiologic studies of lower extremity arteries, at rest and following treadmill stress testing, (i.e., bidirectional Doppler waveform or volume plethysmography recording and analysis at rest with ankle/brachial indices immediately after and at timed intervals following performance of a standardized protocol on a motorized treadmill plus recording of time of onset of claudication or other symptoms, maximal walking time, and time to recovery) complete bilateral study

## Standard Anatomic Coverage

- Arteries of the upper and lower extremities

## Imaging Considerations

- This guideline does not supersede the enrollee's health plan medical policy specific to diagnostic Physiologic Testing for Peripheral Arterial Disease (PAD) of the Upper and Lower Extremities
- This guideline does not address Duplex imaging of the upper and lower extremities (CPT codes 93925, 93926 93930, and 93931) which are covered in a separate guideline document
- For the purposes of the current guideline, Physiological Testing is defined as follows: Evaluation of the peripheral circulation based on measurement of limb blood pressures with pulse volume recordings or Doppler waveforms, or other parameters without utilizing data from direct imaging of the blood vessels
- Selection of the optimal diagnostic imaging for evaluation of peripheral arterial disease should be made within the context of other available modalities (which include Duplex Vascular imaging studies, CT angiography (CTA), Magnetic Resonance angiography (MRA) and contrast angiography) so that the resulting information facilitates patient management decisions and does not merely add a new layer of testing
- Duplicative testing or repeat imaging of the same anatomic area with same or similar technology may be subject to high-level review and may not be medically necessary unless there is a persistent diagnostic problem or there has been a change in clinical status (e.g. deterioration) or there is a medical intervention which warrants interval reassessment
- It is assumed that patients who have cardiovascular disease (established coronary, cerebrovascular disease) and patients at high risk of cardiovascular disease (including patients with diabetes mellitus and chronic kidney disease) will be treated with optimal medical therapy. Therefore, screening for asymptomatic PAD in these populations is unlikely to change management
- Evidence is lacking that treatment of asymptomatic PAD delays the onset of symptomatic PAD
- Although abnormal ankle brachial index (ABI) has been linked to increased incidence of cardiovascular events, treatment of asymptomatic patients with abnormal ABI (who would not otherwise merit treatment) has not been shown to improve outcomes

- ABI has been used in combination with other risk tools (e.g. Framingham Risk Score) and in some cases has led to reclassification of risk (either upwards or downwards). Management of such patients per their reclassified risk level has not been shown to improve outcomes
- It is conventional to report ABI measurements as follows: Noncompressible values defined as greater than 1.40, normal values 1.00 to 1.40, borderline 0.91 to 0.99, and abnormal 0.90 or less
- ABI of 0.9 or less establishes the diagnosis of PAD. Further imaging studies are required only when the patient is being evaluated for revascularization
- Normal resting physiologic studies may be followed by exercise studies at the discretion of the provider. Exercise studies are usually not appropriate when the resting study is abnormal

## Diagnostic Physiologic Testing for PAD of the Lower Extremities

The following diagnostic indications for Physiologic Testing for Peripheral Arterial Disease of the Lower Extremities are accompanied by pre-test considerations as well as supporting clinical data and prerequisite information:

### Asymptomatic patients with suspected PAD

- Screening for asymptomatic PAD has not been shown to affect outcomes and is therefore not considered medically necessary

### Symptomatic patients with suspected PAD

(any one of the following)

- Physiological testing is appropriate for patients with new or worsening exertional limb symptoms (claudication)
- Physiological testing is appropriate for resting limb pain thought to be due to ischemia (because of diminished or absent pulses)
- Physiological testing is appropriate for patients with non-healing ulcers or gangrene of the lower extremities
- Physiological testing is appropriate for patients with infection of the leg or foot with no palpable pulses
- Physiological testing is appropriate for patients with suspected acute limb ischemia (suggested by sudden onset of pain associated with pulselessness, pallor, loss of motor or sensory function)

### Patients with established PAD who have not undergone revascularization

(any one of the following)

- Bilateral ABI should be measured in all patients with newly diagnosed PAD to establish a baseline
- Physiological testing is appropriate for patients with new or worsening exertional limb symptoms (claudication)
- Physiological testing is appropriate for patients with non-healing leg ulcers or gangrene
- Physiological testing is appropriate for patients with infection of the leg or foot with no palpable pulses
- Segmental pressure measurements are appropriate in patients with established PAD to establish the level of disease when intervention is anticipated

### Patients with established PAD who have undergone revascularization

(any one of the following)

- Physiological testing is appropriate for patients with new or worsening exertional limb symptoms (claudication)
- Physiological testing is appropriate for patients with leg ulcers or gangrene
- Physiological testing is appropriate for patients with infection of the leg or foot with no palpable pulses
- A post procedure baseline surveillance (no new or worsening symptoms) study is appropriate (usually performed within 6 months of the revascularization procedure)
- A follow-up surveillance (no new or worsening symptoms) study at 6-12 months following surgical revascularization is appropriate. Note that this guideline is not applicable for surveillance following catheter based revascularization (angioplasty, stent placement etc.)
- An annual follow-up surveillance study (no new or worsening symptoms) starting 1 year after revascularization is appropriate
- Segmental pressure measurements are appropriate in patients with established PAD to establish the level of disease when intervention is anticipated

# Diagnostic Physiologic Testing for PAD of the Upper Extremities

The following diagnostic indications for Physiologic Testing for Peripheral Arterial Disease of the Upper Extremities are accompanied by pre-test considerations as well as supporting clinical data and prerequisite information:

## Asymptomatic patients with suspected PAD

- Screening for asymptomatic PAD has not been shown to affect outcomes and is therefore not considered medically necessary

---

## Symptomatic patients with suspected PAD

(any **one** of the following)

- Physiological testing is appropriate for new or worsening exertional arm or hand symptoms (claudication)
- Physiological testing is appropriate for unilateral cold painful hand
- Physiological testing is appropriate for finger discoloration or ulcer
- Physiological testing is appropriate for suspected positional arterial obstruction
- Physiological testing is appropriate for arm or hand trauma and a suspicion of vascular injury
- Physiological testing is appropriate prior to planned harvest of the radial artery (e.g. for CABG)
- Physiological testing is appropriate in the presence of pulsatile mass or hand ischemia after upper extremity vascular access
- Physiological testing is appropriate in the presence of bruit after upper extremity access for intervention

---

## Patients with established PAD who have not undergone revascularization

(any **one** of the following)

- Physiological testing is appropriate for new or worsening exertional arm or hand symptoms (claudication)
- Physiological testing is appropriate for unilateral cold painful hand
- Physiological testing is appropriate for finger discoloration or ulcer
- Physiological testing is appropriate for arm or hand trauma and a suspicion of vascular injury
- Physiological testing is appropriate prior to planned harvest of the radial artery (e.g. for CABG)
- Physiological testing is appropriate in the presence of pulsatile mass or hand ischemia after upper extremity vascular access
- Physiological testing is appropriate in the presence of bruit, thrill, hematoma or severe pain after upper extremity vascular access
- Segmental pressure measurements are appropriate in patients with established PAD to establish the level of disease when intervention is anticipated

---

## Patients with established PAD who have undergone revascularization

(any **one** of the following)

- Physiological testing is appropriate for new or worsening exertional arm or hand symptoms (claudication)
- Physiological testing is appropriate for patients with arm or hand ulcers or gangrene
- Physiological testing is appropriate for patients with infection of the arm or hand with no palpable pulses
- Following trauma to the revascularized limb with suspected vascular injury
- A post procedure baseline surveillance (no new or worsening symptoms) study is appropriate (usually performed within 6 months of the revascularization procedure)
- A follow-up surveillance (no new or worsening symptoms) study at 6-12 months following surgical revascularization using a vein bypass graft is appropriate. Note that this guideline is not applicable for surveillance following either catheter based revascularization (angioplasty, stent placement etc.) or surgical bypass using a prosthetic graft
- An annual follow-up surveillance study (no new or worsening symptoms) starting 1 year after revascularization is appropriate following surgical revascularization using a vein bypass graft or a prosthetic bypass is appropriate

**Note** *This guideline is not applicable for surveillance following catheter based revascularization (angioplasty, stent placement etc.)*

## Reference/Literature Review

1. American College of Cardiology Foundation (ACCF); American College of Radiology (ACR); American Institute of Ultrasound in Medicine (AIUM); et al. ACCF/ACR/AIUM/ASE/ASN/ICAVL/SCAI/SCCT/SIR/SVM/SVS/SVU 2012 appropriate use criteria for peripheral vascular ultrasound and physiological testing part I: arterial ultrasound and physiological testing: a report of the American College of Cardiology Foundation appropriate use criteria task force, American College of Radiology, American Institute of Ultrasound in Medicine, American Society of Echocardiography, American Society of Nephrology, Intersocietal Commission for the Accreditation of Vascular Laboratories, Society for Cardiovascular Angiography and Interventions, Society of Cardiovascular Computed Tomography, Society for Interventional Radiology, Society for Vascular Medicine, Society for Vascular Surgery, [corrected] and Society for Vascular Ultrasound. *J Am Coll Cardiol*. 2012;60(3):242-276.
2. American College of Cardiology Foundation, American Heart Association, Inc. *ACCF/AHA Pocket Guideline: Management of Patients With Peripheral Artery Disease (Lower Extremity, Renal, Mesenteric, and Abdominal Aortic)*. ACCF/AHA; November 2011.
3. Brott TG, Halperin JL, Abbara S, et al. 2011 ASA/ACCF/AHA/AANN/AANS/ACR/ASNR/CNS/SAIP/SCAI/SIR/SNIS/SVM/SVS guideline on the management of patients with extracranial carotid and vertebral artery disease: a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines, and the American Stroke Association, American Association of Neuroscience Nurses, American Association of Neurological Surgeons, American College of Radiology, American Society of Neuroradiology, Congress of Neurological Surgeons, Society of Atherosclerosis Imaging and Prevention, Society for Cardiovascular Angiography and Interventions, Society of Interventional Radiology, Society of NeuroInterventional Surgery, Society for Vascular Medicine, and Society for Vascular Surgery. *J Am Coll Cardiol*. 2011;57(8):e16-e94.
4. Hirsch AT, Haskal ZJ, Hertzner NR, et al. ACC/AHA 2005 Practice Guidelines for the management of patients with peripheral arterial disease (lower extremity, renal, mesenteric, and abdominal aortic): a collaborative report from the American Association for Vascular Surgery/Society for Vascular Surgery, Society for Cardiovascular Angiography and Interventions, Society for Vascular Medicine and Biology, Society of Interventional Radiology, and the ACC/AHA Task Force on Practice Guidelines (Writing Committee to Develop Guidelines for the Management of Patients With Peripheral Arterial Disease): endorsed by the American Association of Cardiovascular and Pulmonary Rehabilitation; National Heart, Lung, and Blood Institute; Society for Vascular Nursing; TransAtlantic Inter-Society Consensus; and Vascular Disease Foundation. *Circulation*. 2006;113(11):e463-e654.
5. Moyer VA; U.S. Preventive Services Task Force. Screening for peripheral artery disease and cardiovascular disease risk assessment with the ankle-brachial index in adults: U.S. Preventive Services Task Force recommendation statement. *Ann Intern Med*. 2013;159(5):342-348.
6. Norgren L, Hiatt WR, Dormandy JA, et al. Inter-Society Consensus for the Management of Peripheral Arterial Disease (TASC II). *Eur J Vasc Endovasc Surg*. 2007;33 Suppl 1:S1-S75.
7. Rooke TW, Hirsch AT, Misra S, et al. 2011 ACCF/AHA Focused Update of the Guideline for the Management of Patients With Peripheral Artery Disease (updating the 2005 guideline): a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines. *J Am Coll Cardiol*. 2011;58(19):2020-2045.
8. Rooke TW, Hirsch AT, Misra S, et al.; American College of Cardiology Foundation Task Force; American Heart Association Task Force. Management of patients with peripheral artery disease (compilation of 2005 and 2011 ACCF/AHA Guideline Recommendations): a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines. *J Am Coll Cardiol*. 2013;61(14):1555-1570.